

Decomposing Björner's Matrix

Światosław R. Gal*

Wrocław University

<http://www.math.uni.wroc.pl/~sgal/papers/bjo.pdf>

Abstract: We give an alternative proof of a (former) conjecture of Björner stating that the matrix expressing face numbers in terms of g numbers is totally non-negative. We briefly discuss the case of simple flag polytopes.

Let $f(t) = \sum f_i t^i$ denote the f -polynomial of a d -dimensional simple polytope. The coefficients (f_i denotes the number of codimension i faces of the polytope) are called *face numbers* of the polytope. It is a consequence of Dehn-Somerville relations that $f(t)$ can be written as a linear combination of polynomials $u_i(t) = \sum_{q=i}^{d-i} (1+t)^q$. The coefficients of the corresponding expansions are denoted g_i i.e.

$$f(t) = \sum_{i=0}^{\lfloor d/2 \rfloor} g_i u_i(t).$$

In particular there exists a matrix $M_d^{(g)}$ such that

$$\begin{pmatrix} f_0 \\ f_1 \\ \vdots \\ f_d \end{pmatrix} = M_d^{(g)} \begin{pmatrix} g_0 \\ g_1 \\ \vdots \\ g_{\lfloor d/2 \rfloor} \end{pmatrix}.$$

It was conjectured by Björner that $M_d^{(g)}$ is *totally non-negative* i.e. all its minors are non-negative. This was proven by Björklund and Engström [BE].

Björner already proved [Bj] that two-by-two minors of $M_d^{(g)}$ are non-negative and used it to refine lower and upper bounds for simple polytopes.

In this note we present another proof which shows that $M_d^{(g)}$ is totally non-negative.

Since face and g -numbers are related as follows

$$\begin{aligned} \sum f_i t^{d-i} &= \sum_{k=0}^{\lfloor d/2 \rfloor} g_k \sum_{j=k}^{d-k} (1+t)^j \\ &= \sum_{k=0}^{\lfloor d/2 \rfloor} g_k \sum_i \sum_{j=k}^{d-k} \binom{j}{d-i} t^{d-i} \\ &= \sum_i t^{d-i} \sum_{k=0}^{\lfloor d/2 \rfloor} g_k \left(\binom{d-k+1}{d-i+1} - \binom{k}{d-i+1} \right). \end{aligned}$$

* Partially supported by a MNISW grant N N201 541738.

2010 *Mathematics Subject Classification*: 52B05.

Key phrases: g -vector, Björner's Conjecture.

We have

$$f_i = \sum_{k=0}^{\lfloor d/2 \rfloor} M_d^{(g)}(i, k) g_k$$

with

$$M_d^{(g)}(i, k) = \binom{d-k+1}{d-i+1} - \binom{k}{d-i+1}.$$

1. FOUR MATRICES.

Let us define four infinite matrices:

$$\begin{aligned} A_{\pm}(i, j) &= \binom{j+1}{i-j} \mp \binom{j}{i-j-1}, \\ G_+(j, k) &= \frac{2k+1}{2j+1} \binom{k+j}{2j}, \\ G_-(j, k) &= \binom{k+j+1}{2j+1}. \end{aligned}$$

In all the cases we assume that i , j , and k run through natural numbers including zero.

Proposition. *The matrices A_+ , A_- , G_+ , and G_- are totally non-negative.*

Proof: First notice that

$$A_+(i, j) = \binom{j}{i-j}, \quad A_-(i, j) = \frac{i+1}{j+1} \binom{j+1}{i-j}.$$

It is quite standard that

$$\binom{j}{i-j}, \quad \binom{j+1}{i-j}, \quad \binom{k+j}{2j}, \quad \binom{k+j+1}{2j+1}$$

are totally non-negative. For example $\binom{k+j}{2j}$ counts the number of paths from $(-2j, j)$ to $(0, k)$ where only steps $(1, 0)$ or $(0, 1)$ are allowed. (Cf. [FZ, Lemma 1].)

When $X(i, j)$ is totally non-negative, then $a(i)X(i, j)b(j)$ is totally non-negative provided a and b are non-negative sequences. Thus the proof. $\square$

2. DECOMPOSITION OF THE MATRIX $M^{(g)}$

Theorem. Let $\varepsilon = (-)^d$ and $n = \lfloor d/2 \rfloor$. One has

$$M_d^{(g)}(i, k) = \sum_{j=0}^n A_\varepsilon(i, j) G_\varepsilon(n - j, n - k).$$

Remark. Since total non-negativeness is preserved under reversing the order of both indices, and multiplication of such matrices total non-negativity of $M_d^{(g)}$ follows immediately. We believe that the above formula is of independent interest.

Poof of the Theorem: It reduces to:

$$\binom{2n - k + 1}{2n - i + 1} - \binom{k}{2n - i + 1} = \sum_{j=k}^n \binom{j}{i - j} \frac{2n - 2k + 1}{2n - 2j + 1} \binom{2n - (k + j)}{2n - 2j} \quad (1)$$

$$\binom{2n - k + 2}{2n - i + 2} - \binom{k}{2n - i + 2} = \sum_{j=k}^n \frac{i + 1}{j + 1} \binom{j + 1}{i - j} \binom{2n - (k + j) + 1}{2n - 2j + 1} \quad (2)$$

Let us provide a proof of the first identity. The second can be obtained in a very similar way.

Multiply both sides by s^i and sum over $0 \leq i \leq 2n + 1$. Left-hand side of (1) becomes

$$s^{2n+1}(1 + 1/s)^{2n-k+1} - s^{2n+1}(1 + 1/s)^k = (1 + s)^{2n+1} \left(\frac{s}{1 + s} \right)^k - s^{2n+1} \left(\frac{1 + s}{s} \right)^k.$$

The right-hand side of (1) becomes

$$\sum_{j=k}^n (s(s + 1))^j \frac{2n - 2k + 1}{2n - 2j + 1} \binom{2n - (k + j)}{2n - 2j}.$$

Let

$$F_a(z) = \sum_{j=0}^a z^j \frac{2a + 1}{2a - 2j + 1} \binom{2a - j}{2a - 2j}.$$

Then left-hand side of (1) equals to $(s(s + 1))^k F_{n-k}(s(s + 1))$. It is straightforward to check that

$$z(4z + 1)F_a''(z) + a(z - a(4z + 1))F_a'(z) + 2a(2a + 1)F_a(z) = 0.$$

If $G_a(s) = F_a(s(s + 1))$ then

$$s(s + 1)G_a''(s) - 2a(1 + 2s)G_a'(s) + 2a(2a + 1)G_a(s) = 0.$$

The above equation is solved also by $(s+1)^{2a+1}$ and s^{2a+1} . Since $G_a(s)$ and $(s+1)^{2a+1} - s^{2a+1}$ have the same (non-zero) coefficients at s and s^{2a} we conclude that

$$G_a(s) = (s+1)^{2a+1} - s^{2a+1},$$

and, what follows the left-hand side of (1) is equal to

$$(s(s+1))^k G_{n-k}(s) = \left(\frac{s}{1+s}\right)^k (s+1)^{2n+1} - \left(\frac{1+s}{s}\right)^k s^{2n+1}.$$

Which completes the proof. $\square$

Question. The matrices $G_\epsilon(n-j, n-k)$ depend on dimension mildly. And they provide, applied to g -numbers another (natural) expansion of f -polynomial. Does this expansion is combinatorially/geometrically useful?

3. FLAG POLYTOPES AND γ -MATRIX.

If the polytope in question is *flag* (ie. all pairwise intersecting families of faces are centered) an interesting expansion of the f -vector is given in terms of polynomials $v_i(t) = (t+1)^i(t+2)^{d-2i}$, ie.

$$f(t) = \sum \gamma_i v_i(t)$$

(cf [Br,G].)

Unsolved question is whether the coefficients γ_i are non-negative in this case ([G, Conjecture 2.1.7]). As before we can write down the matrix which computes the coefficients of f -polynomial in terms of γ_i :

$$\begin{aligned} \sum f_i t^{d-i} &= \sum_{k=0}^{\lfloor d/2 \rfloor} \gamma_k (1+t)^k (2+t)^{d-2k} \\ &= \sum_{k=0}^{\lfloor d/2 \rfloor} \gamma_k \sum_p \sum_q \binom{k}{p} \binom{d-2k}{q} 2^q t^{k-p+d-2k-q} \\ &= \sum_i t^{d-i} \sum_{k=0}^{\lfloor d/2 \rfloor} \gamma_k \sum_q \binom{k}{i-k-q} \binom{d-2k}{q} 2^q. \end{aligned}$$

Thus we have

$$\begin{pmatrix} f_0 \\ f_1 \\ \vdots \\ f_d \end{pmatrix} = M_d^{(\gamma)} \begin{pmatrix} \gamma_0 \\ \gamma_1 \\ \vdots \\ \gamma_{\lfloor d/2 \rfloor} \end{pmatrix},$$

with

$$M_d^{(\gamma)}(i, k) = \sum_j \binom{k}{i-k-j} \binom{d-2k}{j} 2^j.$$

One can also express g_i by γ_j :

$$g_i = \sum_{0 \leq j \leq i} \left(\binom{n-2j}{i-j} - \binom{n-2j}{i-1-j} \right) \gamma_i = \sum_{0 \leq j \leq i} \binom{n-2j+1}{i-j} \frac{n-2i+1}{n-2j+1} \gamma_i.$$

Since the matrix with entries $\binom{n-2j+1}{i-j}$ is easily seen to be totally non-negative we see that $M_d^{(\gamma)}$ is totally non-negative as $M_d^{(g)}$ is such.

This could be also proven directly by checking that

$$M_d^{(\gamma)}(i, k) = \sum_{j=0}^n A_\varepsilon(i, j) \Gamma(n-j, n-k),$$

where $n = \lfloor d/2 \rfloor$, $\varepsilon = (-)^d$, and

$$\Gamma(j, k) = 4^{k-j} \binom{k}{j}.$$

Question. Can one use non-negativity of (say, two-by-two) minors of $M_d^{(\gamma)}$ to get a (depending on conjectural non-negativity of γ numbers) upper/lower bound for face numbers of flag simple polytopes?

References

- [Bj] A. BJÖRNER, *A comparison theorem for f -vectors of simplicial polytopes*, Pure Appl. Math. Q. **3** (2007), no. 1, part 3, pp. 347–356,
- [BE] M. BJÖRKLUND & A. ENGSTRÖM, *The g -theorem matrices are totally non-negative*, J. Combin. Theory Ser. A, **116** (2009), no. 3, pp. 730–732,
- [Br] P. BRÄNDÉN, *Sign-graded posets, unimodality of WW-polynomials and the Charney-Davis conjecture*, Electron. J. Combin. **11** (2004/06), no. 2, Research Paper 9, (electronic),
- [FZ] S.FOMIN & A.ZELEVINSKY *Total positivity: tests and parametrizations*, Math. Intelligencer **22** (2000), no. 1, pp. 23–33,
- [G] Ś. GAL, *Real root conjecture fails for five- and higher-dimensional spheres*, Discrete Comput. Geom. **34** (2005), no. 2, pp. 269–284.

Mathematical Institute
University of Wrocław
pl. Grunwaldzki 2/4
50-384 Wrocław
Poland
sgal@math.uni.wroc.pl